




SAS-Chicago



Please visit our linked business cards on the [last page](#) of this flyer.

February Meeting

Tuesday, February 13, 2007

The February 2007 meeting will be held at the Angelo's Restorante & Banquets, located at 247 N. York Rd., Elmhurst, IL 60126. See the map on the following page.

Social Hour: 5:30 - 6:30 PM

Dinner: 6:30 PM

Speaker: 7:30 – 9:00 PM

SAS Chicago - Student Night

Time-Resolved Ultraviolet Raman Spectroscopy of Coke Formation

Presentation by Paula M. Allotta

Department of Chemistry and Center for Catalysis and Surface Science,
Northwestern University, Evanston, Illinois 60208

Analyte Detection Using Nanowires Produced by On-Wire Lithography via Confocal Raman Microscopy

Presentation by Lidong Qin

Department of Chemistry
Northwestern University, Evanston, Illinois 60208

Imaging of Surface-Enhanced Raman Scattering “Hotspots” by the Photochemical Reduction of a Self-Assembled Monolayer

Presentation by Matthew A. Young

Department of Chemistry
Northwestern University, Evanston, Illinois 60208

Please make your dinner reservations for the upcoming meeting by email at sas.chicago@bigfoot.com, by using the form on our web page < <http://sas-chicago.org/Online%20Registration%20at%20Angelo.htm> > or by calling Mary Kaplar at 708-449-5767. Leave your name, company affiliation, a telephone number, the number of reservations and your choice of entree. Please call by noon Friday, Feb. 9th, so that proper arrangements can be made with the restaurant. If you can't attend, cancel by Mon. noon: SAS is charged for no-shows.

Entree choices: 6 oz Ribeye Steak, Chicken Florentina Spinach, or Pasta with Garlic and Oil Fried Vegetables.

Dinner Cost Members: \$25

Students and Unemployed Members: \$10
(No Shows may be Billed)

Non-members: \$30

Time-Resolved Ultraviolet Raman Spectroscopy of Coke Formation

Paula M. Allotta¹ and Peter C. Stair^{1,2}

¹*Department of Chemistry and Center for Catalysis and Surface Science,
Northwestern University, Evanston, Illinois 60208*

²*Chemistry Division, Argonne National Laboratory, Argonne, IL 60439*

As with many hydrocarbon conversion catalysts, zeolite H-MFI suffers from deactivation due to carbonaceous buildup (coke) during the methanol-to-gasoline reaction. A traditional method for studying coke formation is to dissolve the zeolite and products and extract the soluble components, which are then studied by GC-MS. This method does not give any information about insoluble coke, and the strong acids and bases used during the extraction process may cause chemistry. Raman spectroscopy has advantages over this GC-MS analysis because measurements can be taken *in-situ*. Visible excitation causes strong fluorescence in coke systems and peaks are often undetectable. By shifting the excitation wavelength to the UV region, fluorescence can be avoided, and peak intensities increase due to frequency dependence.¹ Using UV-Raman spectroscopy, reaction products can be monitored as the reaction is occurring. However, heating and cooling is slow (minutes), and coke formation is too fast for real-time UV-Raman measurements. By introducing an infrared laser as the heat source, heating and cooling are much faster, 10⁻⁶ to 10⁻³ s. After heating, products and intermediates are trapped inside the zeolite pores and unable to react or desorb. Using this pulse-quench sequence, intermediates can be detected by UV-Raman spectroscopy.

(1) Chua, Y. T.; Stair, P. C., *Journal of Catalysis* 2003, 213, 39.

Analyte Detection Using Nanowires Produced by On-Wire Lithography via Confocal Raman Microscopy

Lidong Qin and Chad A Mirkin

Department of Chemistry

Northwestern University, Evanston, Illinois 60208

We have developed a new strategy for designing, fabricating, and imaging Raman hot spots in the context of one-dimensional nanostructures prepared by On-Wire Lithography (OWL).¹⁻³ A combinatorial analysis of these structures provide insight into the electromagnetic mechanism of Surface Enhanced Raman Scattering (SERS). Gold nanodisk arrays with disk thicknesses and interparticle gaps precisely controlled over the 5 nm to many micrometer range were fabricated by OWL and studied by confocal Raman microscopy with methylene blue as a probe molecule. The results demonstrate that disk thickness and gap distance play a crucial role in determining electromagnetic enhancement as determined by local SERS intensities. In contrast to earlier experimental and theoretical work that suggested that the SERS enhancement is largest for junctions with the smallest gap, here we find a non-zero optimum gap size for a specified excitation wavelength that results from the red-shifting of the dipole plasmon wavelength as gap size is decreased. These approaches of using nanowires and Raman spectroscopy open new avenues for detection of analyte and sensing biological and chemical molecules.

(1) Qin, L. D.; Zou, S. L.; Xue, C.; Atkinson, A.; Schatz, G. C.; Mirkin, C. A., *Proceedings of the National Academy of Sciences of the United States of America* 2006, 103, 13300.

(2) Hurst, S. J.; Payne, E. K.; Qin, L. D.; Mirkin, C. A., *Angewandte Chemie-International Edition* 2006, 45, 2672.

(3) Qin, L. D.; Park, S.; Huang, L.; Mirkin, C. A., *Science* 2005, 309, 113.

Imaging of Surface-Enhanced Raman Scattering “Hotspots” by the Photochemical Reduction of a Self-Assembled Monolayer

Matthew A. Young and Richard P. Van Duyne

Department of Chemistry

Northwestern University, Evanston, Illinois 60208


Upon irradiation with green laser light, a self-assembled monolayer of 4-nitrobenzenethiol (NBT) undergoes photochemical reduction to 4-aminobenzenethiol (ABT).¹ It has been shown that this reduction only occurs on surfaces that show surface-enhanced Raman scattering (SERS) activity.² In this study, we have followed the photochemical reduction in real time using SERS and used it to generate near-field images of Ag nanoparticles showing specific areas of extra high SERS activity, or hotspots. Additionally, we have explored the use of this photoconversion for ultra high resolution binary monolayer lithography and have performed some investigations into the mechanism of the photoreduction reaction.

(1) Han, S. W.; Lee, I.; Kim, K. *Langmuir* 2002, 18, 182.

(2) Kim, K.; Lee, I. *Langmuir* 2004, 20, 7351.

DIRECTIONS TO ANGELO’S RESTORANTE & BANQUETS

247 N York St, Elmhurst, IL 60126; Phone: 630-833-2400 - Angelo's Restorante and Banquets is located in Elmhurst less than one block south of North Avenue (IL-64) on York St. on the east side of the street.



Index of Newsletter Sponsors

(Click on card or URL to link to Sponsor's web site)




† **Jeff Kukuk**
Senior Sales Engineer
Inorganic Analysis

† PerkinElmer, Inc.
2000 York Road, Suite 132
Oak Brook, IL 60523
Phone: 800 762-4000 ext. 132
Fax: 630 556-4523
Cell: 630 222-8921
jeff.kukuk@perkinelmer.com
www.perkinelmer.com

www.perkinelmer.com

[email](mailto:jeff.kukuk@perkinelmer.com)

Chip McCauslin
Sales Representative-Atomic Spectroscopy
Analytical Chemist
Chicago Area
Optical Spectroscopy Instruments



Varian Analytical Instruments
2700 Mitchell Drive
Walnut Creek, CA 94598
Phone: 800.926.3000 Ext. 3025
Fax: 925.945.2360
<http://www.varianinc.com>
chip.mccauslin@varianinc.com

www.varianinc.com

[email](mailto:chip.mccauslin@varianinc.com)



For the best in nebulizers, sample introduction components, torches, and spray chambers, call:
1-800-MEINHARD

Meinhard Glass Products

Geoff Coleman, Ph.D., Vice President

700 Corporate Circle, Suite A
Golden, Colorado 80401-5636 USA
Tel: 303.277.9776 Fax: 303.216.2649
gcoleman@meinhard.com www.meinhard.com

www.meinhard.com

[email](mailto:gcoleman@meinhard.com)



† **Steve Bouffard, Ph.D.**
Technical Sales Specialist
FTIR, UV/Vis/NIR, GC, GC/MS
HPLC, DSC, TGA, TMA and DMA

† **PerkinElmer Life and Analytical Science**
2000 York Road, Suite 132
Oak Brook, IL 60523
Phone: (800) 762-4000
Fax: (203) 944-4914
Steven.Bouffard@perkinelmer.com
www.perkinelmer.com

www.perkinelmer.com

[email](mailto:Steven.Bouffard@perkinelmer.com)



5225 Verona Road
Madison, WI
53711-4495

Andrew M. Haefner, Ph.D.
Sales Engineer
Molecular Spectroscopy


(800) 648-5456
(815) 479-0127 fax
(815) 479 0887 direct

Nicolet FT-IR, Raman
FT-NIR, and Microscopy

andrew.haefner@thermo.com www.thermo.com

www.thermo.com

[email](mailto:andrew.haefner@thermo.com)



'Tools for Discoveries of Life'

Rina K. Dukor, Ph.D.
President / CEO

Tel: (847) 487-5500
Fax: (847) 487-5544
Toll Free: (866) BTOOLS1
E-mail: rkdukor@btools.com
Website: www.btools.com

Bio Tools, Inc.
950 N. Rand Rd. Unit 123
Wauconda, IL 60084 USA

www.btools.com

[email](mailto:rkdukor@btools.com)



JERRY SHKOLNIK
Executive Sales Representative
Chicago, IL

TELEDYNE INSTRUMENTS
Leeman Labs
A Teledyne Technologies Company

6 Wentworth Drive
Hudson, New Hampshire 03051
Phone: 847.478.0452 Fax: 847.478.0467
Toll Free: 1.800.634.9942 Cell: 603.493.6176
E-Mail: JShkolnik@Teledyne.com
www.LeemanLabs.com

www.leemanlabs.com

[email](mailto:JShkolnik@Teledyne.com)

Renishaw Inc
5277 Trillium Blvd.
Hoffman Estates, IL 60192
Tel (847) 286-9953 Fax (847) 286-9974
www.renishaw.com



Tim Prusnick
Midwest Sales Manager - Spectroscopy Products Division
Email timothy.prusnick@renishaw.com



www.renishaw.com/spectroscopy

[email](mailto:timothy.prusnick@renishaw.com)

Your company offering a seminar that would be of interest to our membership? Contact Mary Kaplar at mkaplar@bodycoteusa.com to find out how the Chicago SAS can help spread the word.